## (19) World Intellectual Property Organization International Bureau





(43) International Publication Date 29 December 2004 (29.12.2004)

**PCT** 

## (10) International Publication Number WO 2004/113564 A1

(51) International Patent Classification<sup>7</sup>: C07H 21/00, A61K 31/712

C12Q 1/68,

(21) International Application Number:

PCT/EP2004/006534

(22) International Filing Date: 17 June 2004 (17.06.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data: 10328813.9

20 June 2003 (20.06.2003) DE

- (71) Applicant (for all designated States except US): EPIGE-NOMICS AG [DE/DE]; Kastanienallee 24, 10435 Berlin (DE).
- (72) Inventor; and
- (75) Inventor/Applicant (for US only): SCHUSTER, Matthias [DE/DE]; Kastanienallee 107, 13158 Berlin (DE).
- (74) Agent: SCHUBERT, Klemens; Neue Promenade 5, 10178 Berlin (DE).


- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

## Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

[Continued on next page]

(54) Title: METHOD FOR INVESTIGATING CYTOSINE METHYLATION IN DNA SEQUENCES BY MEANS OF TRIPLEX-FORMING OLIGOMERS



(57) Abstract: A method is described for the investigation of cytosine methylation in DNA sequences. Triplex-forming oligomers are utilized, which preferably form triplex structures at positions where cytosine unmethylated at position 5 is present. The triplexes block the transcription, replication and amplification of the DNA. In particular, peptide nucleic acid oligomers with modified nucleobases can be used as triplex-forming oligomers.

## 

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.